



## Modeling eco-hydrological impacts of stream temperature changes in a catchment

Loinaz, Maria Christina; Bauer-Gottwein, Peter; Butts, Michael

*Publication date:*  
2011

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Loinaz, M. C. (Author), Bauer-Gottwein, P. (Author), & Butts, M. (Author). (2011). Modeling eco-hydrological impacts of stream temperature changes in a catchment. Sound/Visual production (digital)

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Modeling eco-hydrological impacts of stream temperature changes in a catchment



HydroEco Conference, Vienna, May 2-5 2011

Maria Loinaz, Technical University of Denmark

Peter Bauer-Gottwein, Technical University of Denmark

Michael Butts, DHI

# Research Objectives


1. Develop **integrated hydro-ecological models** to relate environmental stressors in a catchment to an indicator of ecosystem status.
2. To **quantify changes in ecosystem status** caused by changes in catchment management.



# Stream Temperature: a critical element in fish ecology

- Short period exposure: death
  - Lethal temperatures (LT50)
- Long period exposure: physiological responses
  - Metabolism
  - Food intake
- Important factors:
  - Food availability
  - Habitat conditions
    - Oxygen, pollution
    - Physical stream conditions

Effects of temperature on salmonids





# Catchment factors that impact stream temperature

- Land use/water use impact on flows
  - Groundwater
  - Tributaries
  - Agricultural drains
- Geomorphology
  - Channel width-to-depth increase (bank erosion, accumulation of sediments)
  - Channelization
  - Ponding
- Bank Vegetation (shade)







# Silver Creek Basin, Idaho

Silver Creek is a spring-fed system abundant in wildlife and also a valuable trout habitat.

The climate is of a semi-desert valley with low precipitation and high evaporation.


Approximately 60% of the lower valley is cultivated, of which 80% is irrigated.




# Modeling approach

## 3 scales

1. Surface-Groundwater Model (Wood River Aquifer)
2. Flow-Temperature Model (Silver Creek Basin)
3. Fish Model (Stream Segments)



# Flow Model (MIKE SHE)



Spatially distributed model, 300m resolution

3 geologic layers, 62 soil types, 20 land use types, distributed irrigation system

207 km total stream and canal length, ~200 cross-sections



# Stream Temperature Model (MIKE 11-Ecolab)

$$T = \frac{H}{\rho C_p V} \quad \rho C_p = \text{heat capacity (J/m}^3\text{°C)}$$

## 1D Surface water heat transport:

$$\frac{\partial T}{\partial t} = -\frac{1}{A} \frac{\partial (QT)}{\partial x} + \frac{1}{A} \frac{\partial}{\partial x} \left( DA \frac{\partial T}{\partial x} \right) + \frac{H}{\rho C_p d}$$

Q = flow


A = surface area

D = longitudinal dispersion

d = water depth

H = atmosphere and sediment heat flux

$$\text{Net heat flux into the water (H)} = H_S + H_A - H_C - H_E - H_{LW} + H_G$$



$H_S$  = net solar radiation

$H_A$  = net atmospheric long wave radiation

$H_C$  = sensible heat flux between water and atmosphere ( $T_w$ )

$H_E$  = evaporation/condensation ( $T_w$ )

$H_{LW}$  = outgoing long wave radiation from water ( $T_w$ )

$H_G$  = groundwater heat flux (ag drains + baseflow)

# Bioenergetics: Fish growth model

- Based on energy balance:

Consumption = Metabolism + Wastes + Growth





- All processes are size and temperature dependent

- Growth model:  $\Delta w_i = g_i \cdot w_i$

$\Delta w_i$  = current day growth (grams/day)


$g_i$  = specific growth rate (g/g·day)

$w_i$  = weight at day  $i$  (grams)




# Ecological Suitability Indicators

- Fish growth in under certain conditions relation to growth under natural conditions at maturity age:  $\text{weight}_{\text{scenario}} / \text{weight}_{\text{natural}}$
- Reproduction capability and time it takes to reach maturity





# Flow Calibration

## Picabo St @ Silver Creek




## Stanton Crossing St @ Big Wood River






# Temperature Calibration

Simulated Temperature @ Picabo St. [deg C] —  
 Measured Temperature @ Picabo St. [deg C] ○



$$ME_i = \bar{E}_i = \frac{\sum (Obs_{i,t} - Calc_{i,t})}{n}$$



# Model Scenarios

1. Natural System (NS): Replace agriculture and urban with natural areas, eliminate all canals and irrigation system
2. Crop change (ALF2BAR): alfalfa to barley
3. Land use change (ALF2GRS): alfalfa to grass
4. Bank vegetation (VH): increase bank vegetation
5. Geomorphology (WOKP): remove Kilpatrick dam and pond, replace with narrower cross-sections and natural profile.






# Temperature Results



Average Temperature Change

Seg	VH	WOKP	ALF2BAR	ALF2GRS	NS
1	0.0	0.0	0.1	0.6	-0.5
2	0.0	0.0	-0.2	-0.6	0.0
3	0.0	0.0	0.0	-0.3	-1.7
4	0.0	0.0	-0.1	-0.1	-1.2
5	0.0	0.1	-0.2	-0.6	-1.5
6	0.0	0.2	-0.1	-0.5	-1.6
7	0.0	0.1	-0.1	-0.3	-1.6
8	0.0	0.0	0.0	-0.2	-0.8
9	0.0	0.0	0.1	0.2	-1.0
10	0.0	0.0	0.0	0.0	-3.2
12	0.0	0.0	0.1	0.2	-0.6
13	0.0	0.0	0.0	-0.1	-1.0
15	0.0	0.0	-0.3	-0.9	-1.6
16	0.0	0.0	0.0	-0.2	-1.4
17	0.0	0.0	0.0	0.0	-1.3

Average Summer Water Temperature  
(deg C)



# Ecological Model Results



	F3	T <sub>M</sub> (d)
RT NS	2534	395
RT EC	1573	480
BT NS	2575	493
BT EC	1875	659

F3 = Potential fecundity at age 3

T<sub>M</sub> = time to reach potential maturity (200 mm)



# Conclusions

- All model scenarios cause a change stream temperatures from current conditions, but only scenario 3 (changing alfalfa crops for grass) was significant decrease in average summer temperatures. A combination of different restorations strategies needs to be further explored.
- Ecological model results imply that the current system is at approximately half of its potential.
- The bioenergetics model should also include habitat degradation conditions (p-value).
- Deviation from natural conditions seem to have less of an impact on brown trout than rainbow trout, which may explain the increasing numbers of brown trout in relation to rainbow trout.

# Thank you

